

FRUIT MAX

Reg No: K11107 Act 36 of 1947
FERTILISER GROUP 1

PRODUCT DESCRIPTION:

A Liquid Phosphorus & Potassium Containing Foliar Feed

ACTIVE INGREDIENTS:

Potassium (K)	230.00 g/kg	355.50 g/L
Phosphorus (P)	90.00 g/kg	139.50 g/L
Specific Gravity (SG) at 20°C	1.55	
pH (10% Solution)	9.0	

**NET MASS
VOLUME 20 L**

REGISTRATION HOLDER

Barinor New Business Ventures 2 (Pty) Ltd
CK 2002/015113/07
51 Verdun Road, Kya Sands, 2196
Tel: +21 913 3512



**EMERGENCY CONTACT:
0861 Metson (638 766)**

Batch No: _____
Packed Date: _____

UN Number: Not Regulated



HAZARD STATEMENTS:

- Causes skin irritation.
- Causes serious eye damage.



WARNING:

Keep away from food and feeds. Store in a cool dry place out of reach from children and/or uninformed persons and animals.

PRECAUTIONARY STATEMENTS:

- Poisonous when swallowed.
- Wear protective eye protection.
- Do not eat, drink, or smoke when mixing or using this product.
- Do not inhale dust or vapour.
- Remove contaminated clothing immediately.
- IF ON SKIN: wash with plenty of soap and water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and continue rinsing.
- In case of poisoning: CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM/HER
- Store in a well-ventilated area.
- Keep container tightly closed.
- Destroy empty container by perforating and flattening and dispose of container in a safe way.
- Store away from food and feedstuffs.
- Prevent contamination of food, feedstuffs, drinking water, eating utensils.
- Keep out of reach of children.

DIRECTIONS FOR USE: Use only as directed.

- For best results, apply *FruitMax* as part of a balanced nutritional programme.
- *FruitMax* is compatible with Nitrogen and micro-element containing products. If in doubt, it is advisable to check compatibility by mixing a small quantity first or consult your nearest representative.
- Avoid applying *FruitMax* with Ca and Mg containing products irrespective of the form which it is sold in, i.e. nitrates, sulphates, carbohydrate chelates, amino acid chelates, oxides, carbonates, chlorides, due to incompatibility.
- Avoid mixing *FruitMax* with highly alkaline materials such as Bordeaux Mixture and Lime Sulphur.
- *FruitMax* is suitable for all types of spraying and watering systems.

- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.
- Spray early morning or later afternoon.
- Do not apply during stress periods.
- Keep container tightly closed.
- Recommended storage temperature is below 40°C. Avoid freezing.
- Do not store diluted product.
- Do not apply as a concentrate. Always dilute *FRUITMAX* with water before use.
- Follow the standard mixing procedures.

APPLICATION DETAILS:

Shake well before use

CROPS	APPLICATION RATE	MAX APPLICATION RATE	APPLICATION DETAILS
CITRUS	250 - 500 ml per 100 L water	12 L/ha	Apply twice starting at 100% full bloom. Repeat two weeks after fruit set. Use lower rate on soft Citrus. During fruit colouring & enlargement stage: Apply three times after petal fall. Do not apply within 30 days before harvest. Apply early morning or late afternoon.
BEAN CROPS	200 ml / 100 L water	4 L/ha	2 applications, 1 st application before at flowering and 2 nd application at pod formation.
DECIDUOUS & STONE FRUIT	500 ml per 100 L water	12 L/ha	Apply twice or three times at 21day intervals after petal fall. It is recommended to apply post-harvest as a foliar spray to the tree. During fruit colouring & enlargement stage: Two to three applications at 14day intervals, starting 6 weeks before harvest.
HOPS	250 - 500 ml per 100 L water	12 L/ha	Apply before crop's reproductive growth stage.
NUT TREES	500 ml per 100 L water	12 L/ha	Apply twice starting at full bloom. Repeat two weeks after fruit set.
SUB-TROPICAL FRUIT	500 ml per 100 L water	12 L/ha	Apply twice between bud-break & onset of flowering (2/3 of growth flush completed). Recommended to apply post-harvest as a foliar spray to the tree. During fruit colouring & enlargement stage: Apply three times, starting 2 weeks after petal fall. Do not apply within 30 days before harvest. Apply early morning or late afternoon.
TABLE GRAPES	500 - 750 ml per 100 L water	12 L/ha	Apply at veraison & 7 days later. Recommended to apply post-harvest as a foliar spray to the vines. During fruit colouring & enlargement stage: Two applications at 14-day intervals, starting 6 weeks before harvest.
WINE GRAPES	500 - 750 ml per 100 L water	12 L/ha	Apply at veraison & 7 days later. Recommended to apply post-harvest as a foliar spray to the vines.
VEGETABLES	250 - 500 ml per 100 L water	5 L/ha	Apply before crop's reproductive growth stage.

DISCLAIMERS / CONDITIONS OF SALE:

This product is manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of the manufacturer. Before using, the user should determine the suitability of the product for his/ her intended use and the user assumes the risk and liability in connection therewith.