

# Ascokelp®



Reg No: M216 Act36 of 1947  
FERTILISER GROUP 3

## PRODUCT DESCRIPTION:

**ASCOKELP®** is a seaweed foliar spray

## ACTIVE INGREDIENTS:

Extract from Brown Seaweed (*Ascophyllum nodosum*)

Specific Gravity (SG) @ 20°C 1.01

pH (10% Solution) 6.0 - 7.0

NET MASS  
VOLUME **20 L**

**REGISTRATION HOLDER:** Metson Manufacturing CC  
1999/059914/23  
51 Verdun Road, Kya Sands, 2196  
Tel: + 27 11 792 8968



**EMERGENCY CONTACT:** 0861 Metson (638 766)



**DANGER**



Batch No:  
Packed Date:

UN Number: Not Regulated

## HAZARD STATEMENTS:

- May cause cancer

## PRECAUTIONARY STATEMENTS:

- Poisonous when swallowed Wear protective eye protection
- Do not eat, drink, or smoke when mixing or using this product Do not inhale dust
- Remove contaminated clothing immediately
- IF ON SKIN: wash with plenty of soap and water
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and continue rinsing
- In case of poisoning: CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM/HER.

## STORAGE CONDITIONS:

- Store in a well-ventilated area. Keep container tightly closed.
- Destroy empty container by perforating and flattening and dispose of container in a safe way.
- Store away from food and feedstuffs.
- Prevent contamination of food, feedstuffs, drinking water, eating utensils.
- Keep out of reach of children

## WARNING:

- Keep away from food and feeds. Store in a cool dry place out of reach from children and/or uninformed persons and animals.

## CLAIMS:

- Improves crop growth
- Improves crop yield

## DIRECTIONS FOR USE:

 Use only as directed.

- **ASCOKELP®** is suitable for use with a knapsack, a boom/tractor sprayer, drip irrigation, pivots and aerial application.
- **ASCOKELP®** should be applied after stress periods in order to assist the recovery of the plant.
- Keep container tightly closed.
- Recommended storage temperature below 40°C.
- Avoid freezing.
- Do not store diluted product.
- Always dilute **ASCOKELP®** with water before using.

## APPLICATION DETAILS

☰ Shake well before use

CROP	APPLICATION RATE	APPLICATION DETAIL
Dry beans, Groundnuts & Soya	2 L/ha	<b>First application:</b> Dry beans: V1 Groundnuts: V1 Soya: VC-V1
	2 L/ha	<b>Second application:</b> Dry beans: R1-R3 Groundnuts: R1-R3 Soya: V3-R1
Grapes (Sultana)	3 L/ha	<b>Three foliar applications:</b> 1. 3-4 mm berry size 2. 5-6 mm berry size 3. 9-10 mm berry size  Apply with 1000 L water / ha
Maize	2 L/ha	In furrow at plant &/or postemergence at 4-8 leaf stage.
Wheat	2 L/ha	4-8 leaf stage.
	2 L/ha	Flag leaf stage.

For more information contact your nearest  
Metson Regional Manager

## DISCLAIMERS / CONDITIONS OF SALE:

This product is manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of the manufacturer. Before using, the user should determine the suitability of the product for his/ her intended use and the user assumes the risk and liability in connection therewith.