

# Maxonia™

Reg No: M373 Act 36 of 1947

Patent No.: 2021/10366

FERTILISER GROUP 3

## PRODUCT DESCRIPTION:

A Liquid Bio-Stimulant Containing Seaweed.

## ACTIVE INGREDIENTS:

Phosphorus (P)	4.60 g/kg	4.74 g/L
Potassium (K)	16.80 g/kg	17.30 g/L
Extract from Brown Seaweed ( <i>Ecklonia maxima</i> )		
Specific Gravity (SG) @ 20°C	1.03	
pH (10% Solution)	10.0 - 11.0	

NET MASS  
VOLUME **20 L**

REGISTRATION HOLDER: Metson Biologicals (Pty) Ltd  
CK 2010/006380/06  
51 Verdun Road, Kya Sands, 2196  
Tel: +27 11 792 8968

EMERGENCY CONTACT: 0861 638 706



WARNING  
DANGER



Batch No:  
Packed Date:

UN Number: Not Regulated

WARNING:   
Keep away from food and feeds.  
Store in a cool dry place out of  
reach from children and/or  
uninformed persons and animals.

## HAZARD STATEMENTS:

- Harmful if swallowed
- Causes severe skin burns and eye damage

## PRECAUTIONARY STATEMENTS:

- Poisonous when swallowed
- Wear protective eye-wear
- Do not eat, drink, or smoke when mixing or using this product
- Do not inhale dust or vapour
- Remove contaminated clothing immediately
- IF ON SKIN: wash with plenty of soap and water
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and continue rinsing
- In case of poisoning: CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM/HER.

## STORAGE CONDITIONS :

- Store in a well-ventilated area. Keep container tightly closed.
- Destroy empty container by perforating, flattening and dispose of container in a safe way.
- Store away from food and feedstuffs.
- Prevent contamination of food, feedstuffs, drinking water; eating utensils.
- Keep out of reach of children.
- Keep from freezing and store in a cool place.



## Claims

- Increased Root mass
- Increased Shoot mass
- Increases Yields

## DIRECTIONS FOR USE: Use only as directed.

Refer to Application Table below.

- Shake well before use.
- *Maxonia™* is suitable for use with a knapsack, a boom / tractor sprayer, drip irrigation, pivots and aerial application.
- Always dilute *Maxonia™* with water before using.
- Do not apply undiluted.

## APPLICATION DETAILS

 Shake well before use

Crops	Application Rate	Application Details
<b>Row crops: Cereal</b> Maize, Millet, Sweetcorn, Sorghum, Wheat, Barley, Oat, Rye, Triticale, and all other cereal crops.	2 L/ha	<b>First application:</b> In-furrow.
<b>Legume and Oilseed crops:</b> Dry bean, Green Bean, Pea, Soy, Lentil, Lupin, Clover, Alfalfa, Sunflower, Canola, Rapeseed, Groundnut, Cotton, Tobacco, and all other legume and oilseed crops.		<b>Second application:</b> Post-emergence at 4-8 leaf stage.
<b>Citrus</b>	2 - 4 L/ha	<b>Growth stages:</b> White Bud Full Bloom Petal Fall

For more information contact your nearest Metson Regional Manager.

## DISCLAIMERS / CONDITIONS OF SALE:

This product is manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of the manufacturer. Before using, the user should determine the suitability of the product for his/ her intended use and the user assumes the risk and liability in connection therewith.