



BioMax

Reg No: M468 Act 36 of 1947
FERTILISER GROUP 3

PRODUCT DESCRIPTION:

A liquid seaweed extract.

ACTIVE INGREDIENTS:

Phosphorus (P) 4.60 g/kg 4.74 g/L
 Potassium (K) 16.80 g/kg 17.3 g/L
 Extracts from Brown Seaweed (*Ecklonia maxima*)
 Specific gravity (SG) at 20°C 1.03
 pH (10% Solution) 10.0 - 11.0

NET MASS **20 L**
VOLUME

REGISTRATION HOLDER:

Metson Biologicals
 CK 2010/006380/06
 P O Box 1005, Cresta, 2118
 Tel: +27 11 792 8968



EMERGENCY CONTACT: 0861 Metson (638 766)



**WARNING
 DANGER**



Batch No:
 Packed Date:

UN Number: Not regulated

WARNING:
 Keep away from food and feeds. Store in a cool dry place out of reach from children and/or uninformed persons and animals.

HAZARD STATEMENT:

- Harmful if swallowed.
- Causes severe skin burns and eye damage.

PRECAUTIONARY STATEMENTS:

- Poisonous when swallowed.
- Wear protective eye-wear.
- Do not eat, drink, or smoke when mixing or using this product.
- Do not inhale dust or vapour.
- Remove contaminated clothing immediately.
- IF ON SKIN: Wash with plenty of soap and water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and continue rinsing.
- In case of poisoning: CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM/HER.

STORAGE CONDITIONS:

- Store in a well-ventilated area.
- Keep container tightly closed.
- Destroy empty container by perforating, flattening and dispose of container in a safe way.
- Store away from food or feedstuffs.
- Prevent contamination of food, feedstuffs, drinking water, eating utensils.
- Keep out of reach of children.
- Keep from freezing and store in a cool place.

CLAIMS

- Increased root mass.
- Increased shoot mass.
- Increases yields.

DIRECTIONS FOR USE: (Use only as directed)

Refer to application label below.

- **BIOMAX** is suitable for use with a knapsack, a boom / tractor sprayer, drip irrigation, pivots and aerial application.
- Always dilute **BIOMAX** with water before using.

Shake well before use.

APPLICATION DETAILS:

CROPS	APPLICATION RATE	APPLICATION DETAILS
Legume crops: Dry bean, Green Bean, Pea, Lentil, Lupin, Clover, Alfalfa, Sunflower, Groundnut.	1 - 2 L/ha	Second application: Post-emergence at 4-8 leaf stage.
Row crops: grain Maize, Millet, Sweetcorn, Sorghum.		First application: In-furrow.
Row crops: small grain Wheat, Barley, Oat, Rye, Triticale, and all other cereal crops.		First application: In-furrow.
Other crops: Canola, Cotton, Rapeseed, Tobacco.		Second application: Post-emergence at 4-8 leaf stage.

For more information contact your nearest Metson Regional Manager

DISCLAIMER / CONDITIONS OF SALE:

This product is manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of the manufacturer. Before using, the user should determine the suitability of the product for his/ her intended use and the user assumes the risk and liability in connection therewith.