

# Bambino A2Zn

FERTILIZER GROUP 1 Reg No. K10300, Act 36 of 1947 (RSA)

## 5 kg



**DANGER**

F407005ILT001RSAV1

## BAMBINO A2ZN IS A WATER SOLUBLE POWDER PLANT NUTRIENT.

### COMPOSITION:

**Nitrogen (N)** 185 g/kg, **Phosphorus (P)** 64 g/kg, **Potassium (K)** 40 g/kg, **Magnesium (Mg)** 20 g/kg, **Zinc (Zn)** 42 g/kg, **Manganese (Mn)** 5 g/kg, **Iron (Fe)** 3 g/kg, **Copper (Cu)** 3 g/kg, **Boron (B)** 8 g/kg

### DIRECTIONS FOR USE: Use only as directed.

Apply **Bambino A2Zn** as high volume foliar spray either in the early morning or late afternoon. Do not spray when temperatures exceed 30 °C or onto plants undergoing moisture stress as this can result in reduced uptake or leaf burn. Never apply **Bambino A2Zn** with or within 30 days after application of copper containing fungicide, phosphates and/ or phosphonates. Do not use in combination with highly alkaline material such as Bordeaux Mixtures and Lime Sulphur. The use of a suitable pH buffer is recommended. It is advisable to do a miscibility test prior to mixing **Bambino A2Zn** with other products. Consult with a qualified agronomist for crop specific information. **Bambino A2Zn** should preferably be applied based on recommendations derived from soil and/or foliar analysis.

**RECOMMENDATIONS:** Ensure not to exceed the maximum rate per hectare. Do not exceed the concentration of 2% (2 kg/100 L) when mixing with other agrochemicals in a spray tank.

CROP	RATE	APPLICATION DETAILS
Grapes (Table or Wine) Berries and Shrubs	1 kg/100 L	Start application when sufficient leaves are present. Repeat application at 14 day intervals until sufficient growth is obtained.
Young Non-Bearing Fruit Trees Nursery Trees	1 kg/100 L	Deciduous Trees: Start application when sufficient leaves are present. Repeat application at 14 day intervals. Non-deciduous trees: Start application 14 days post transplanting. Repeat application at 14 day intervals. Apply 4 – 6 applications, or until sufficient growth has been obtained.
Grains Row Crops	3 – 5 kg/ha	Start application at 2 - 3 leaf stage (1 – 3 applications recommended).
Pastures	3 – 5 kg/ha	Apply after every grazing or cutting.

For optimum results **Bambino A2Zn** should be used in a program and in combination with other balanced fertilizers and foliar feeds. Lower rates may be applied for maintenance and higher rates to correct deficiencies.

### RELEVANT SUBSTANCES:

CHEMICAL NAME	CAS No.	EC No.	Conc. %w/w
Disodium Octaborate	12280-03-4	None	<10
EDTA Copper	14025-15-1	None	<10
Zinc Oxide	1314-13-2	None	<10

**DISCLAIMERS:** The information on this sheet is not a specification; it does not guarantee specific properties. The information is intended to provide general guidance as to health and safety based upon our knowledge of the handling, storage and use of the product. It is not applicable to unusual or non-standard uses of the product nor where instructions or recommendations are not followed. All information is given in good faith but without guarantee in respect of accuracy, and no responsibility is accepted for errors and omissions or the consequence thereof.

**HAZARD STATEMENT:** Causes serious eye irritation. May damage fertility. May damage unborn child. Toxic to aquatic life. Toxic to aquatic life with long lasting effects.

**PRECAUTIONARY STATEMENTS:** Wear impervious rubber gloves and chemical safety goggles.  
**IF ON SKIN:** Wash with plenty of water and soap. **IF SKIN IRRITATION OR RASH OCCURS:** Get medical advice. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing. Collect spillage. Dispose of contents/container in accordance with local regulations.

### TRANSPORT INFORMATION

CLASS	PACKAGING GROUP	UN NUMBER	PROPER SHIPPING NAME (PSN)
None	None	None	None

Griffon Poison Centre: +27 (82) 446 8946

SCAN ME/VIEW SDS



**DANGER**

B407005ILT001RSAV1