

BitterPil

REGISTRATION HOLDER:
In Line Trading 112 (Pty) Ltd.
(Co. Reg. No. 2003/008663/07)
Cedar Lake Industrial Estate,
C/O M57 & Porcelain Roads,
Olifantsfontein, 1666, Gauteng, RSA
Tel: +27 (11) 392 4072
Griffon Poison Centre: +27 (82) 446 8946



FERTILIZER GROUP 2 Reg No. K11688, Act 36 of 1947

A FORMULATED PRODUCT FOR THE TREATMENT OF CALCIUM-RELATED DEFICIENCIES IN FRUIT AND OTHER CROPS.

COMPOSITION:
Calcium (Ca) 119 g/kg
SG @ 20°C = 1.29



WARNING

HAZARD STATEMENTS: May be harmful if swallowed. Causes skin irritation. Combustible liquid. Causes serious eye irritation.

PRECAUTIONARY STATEMENTS: Wear impervious rubber gloves and chemical safety goggles. Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Wash hands thoroughly after handling. **IF ON SKIN:** Wash with plenty of water and soap. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing. Collect spillage. Dispose of contents/container in accordance with local regulations.

RELEVANT SUBSTANCES:

CHEMICAL NAME	CAS NO.	EC NO.	CONC. %W/W
Calcium Chloride Flakes	10043-52-4	233-140-8	30-60
Acetic Acid	64-19-7	200-580-7	<10

DIRECTIONS FOR USE: Use only as directed.

COMPATIBILITY: It is advisable to do a miscibility test prior to mixing of BitterPil with other products. Do not apply with any products that contain Sulphates.

MIXING: Shake well prior to opening. Always mix BitterPil with ample quantities of water. Never apply BitterPil undiluted. Follow standard tank mixing procedures.

STORAGE: Ensure BitterPil does not freeze. Store in a cool dry place. Never store in direct sunlight for prolonged periods.

APPLICATION: BitterPil is recommended for the use in conjunction with a complete fertiliser program based on soil or tissue analysis. Should preferably be applied in early morning or late afternoon. Never apply BitterPil to crops undergoing drought stress.

GENERAL: Consult with a qualified agronomist for crop specific information. BitterPil should preferably be applied based on recommendations derived from soil and/or foliar analysis.

APPLICATION TABLE

CROP	RATE/Ha	RATE/100 L
Beans: Alfalfa, Green, Dry, Lupine, Peanut, Soya	2 L – 5 L	250 ml – 1 L
Bulb Vegetables: Garlic, Onion	5 L – 10 L	250 ml – 1 L
Bush & Creeping Vegetables: Cucurbits, Pepper, Strawberry, Tomato	5 L – 10 L	250 ml – 1 L
Leaf Vegetables: Asparagus, Celery, Cruciferous, Lettuce, Spinach	2 L – 5 L	250 ml – 500 ml
Nut Trees: Cashew, Macadamia, Pecan, Pistachio	5 L – 20 L	250 ml – 1 L
Oil Crops: Canola, Sunflower	2 L – 5 L	250 ml – 1 L
Ornamentals: Flowers, Lawns	2 L – 5 L	250 ml – 500 ml
Pome Trees: Apple, Pear, Quince	5 L – 20 L	250 ml – 1 L
Root Vegetables: Beetroot, Carrot, Chicory, Ginger, Turnip	5 L – 10 L	250 ml – 1 L
Shrubs & Other Trees: Blue & Young Berries, Coffee, Cotton, Olive, Pineapple, Pomegranate, Tobacco	5 L – 15 L	250 ml – 500 ml
Stone Trees: Apricot, Cherry, Nectarine, Peach, Plum	5 L – 20 L	250 ml – 1 L
Subtropical Trees: Avocado, Citrus, Guava, Litchi, Mango	5 L – 15 L	150 ml – 500 ml
Sugar Crops: Sugar Beet, Sugarcane	5 L – 10 L	250 ml – 1 L
Summer Grains: Maize, Silage, Sorghum, Sweet corn	1 L – 5 L	250 ml – 1 L
Tropical Trees: Banana, Papaya	5 L – 15 L	250 ml – 500 ml
Tuber Vegetables: Onion, Potato, Yam	5 L – 10 L	250 ml – 1 L
Vines: Granadilla, Hops, Kiwi, Raisin Grape, Table Grape, Wine Grape	5 L – 15 L	250 ml – 500 ml
Winter Grains: Barley, Oats, Wheat	2 L – 5 L	250 ml – 1 L

DISCLAIMERS: This product is manufactured and tested to specific standards and is sold in good faith, however, no guarantee is given as conditions and methods under which they are used are under the control of the manufacturer. Before using, the user should determine the suitability of the product for their intended use and the user assumes the risk and liability in connection therewith.

SEE PACK FOR BATCH DETAILS

20 L



SHAKE WELL BEFORE USE